

# Structural susceptibility and separation of time scales in the van der Pol oscillator

Ricky Chachra, Mark K. Transtrum, and James P. Sethna\*

*Department of Physics, Laboratory of Atomic and Solid State Physics, Cornell University, Ithaca, New York 14853, USA*

(Received 23 December 2011; revised manuscript received 16 April 2012; published 31 August 2012)

We use an extension of the van der Pol oscillator as an example of a system with multiple time scales to study the *susceptibility* of its trajectory to polynomial perturbations in the dynamics. A striking feature of many nonlinear, multiparameter models is an apparently inherent insensitivity to large-magnitude variations in certain linear combinations of parameters. This phenomenon of “sloppiness” is quantified by calculating the eigenvalues of the Hessian matrix of the least-squares cost function. These typically span many orders of magnitude. The van der Pol system is no exception: Perturbations in its dynamics show that most directions in parameter space weakly affect the limit cycle, whereas only a few directions are stiff. With this study, we show that separating the time scales in the van der Pol system leads to a further separation of eigenvalues. Parameter combinations which perturb the slow manifold are stiffer and those which solely affect the jumps in the dynamics are sloppier.

DOI: [10.1103/PhysRevE.86.026712](https://doi.org/10.1103/PhysRevE.86.026712)

PACS number(s): 02.60.Ed

## I. INTRODUCTION

In this paper, we analyze the sensitivity of a multiple-time-scale dynamical system to perturbative changes in its evolution laws. Rather than utilizing the traditional means of examining the *structural stability* for probing qualitative changes to the attractor as a response to perturbations, we study the *structural susceptibility* for quantifying the effects of the perturbations on the time series [1]. More specifically, we ask how sensitive is the dynamical system  $dz/dt = \mathbf{f}(\mathbf{z})$  to infinitesimal changes of the form  $d\mathbf{z}/dt = \mathbf{f}(\mathbf{z}) + \mathbf{a} \cdot \mathbf{g}(\mathbf{z})$  for a family of perturbations  $\mathbf{g}(\mathbf{z})$  when the parameters  $\mathbf{a} \rightarrow \mathbf{0}$ .

This paper introduces the concept of “structural susceptibility” in dynamical systems, and is an outgrowth of our group’s previous work on “sloppiness” in multiparameter systems wherein we have found that data fitting in a number of nonlinear, multiparameter models is only sensitive to a few directions in parameter space at the best fit [2–4]. The key difference between studying sloppiness and structural susceptibilities is that in the former, the parameters are intrinsic to the system, i.e., there are no externally introduced changes in their evolution laws. Nonetheless, the methodology we have developed for studying sloppy models is also suited for studying structural susceptibilities of dynamical systems. Our approach cleanly isolates and ranks the directions in parameter space in order of relevance to observed behavior, and has previously led us to suggest improvements in experimental design [5], extract falsifiable predictions from experiments [6], and develop faster minimization algorithms [7]. Others have developed our ideas to suggest further improvements in experimental design [8] and parameter estimation [9], to quantify robustness to parameter variations [10], and to set confidence regions for predictions in multiscale models [11]. In this paper, we bring similar ideas together to analyze sensitivities of time series to perturbations in dynamical systems.

We demonstrate the utility of our approach with application to a dynamical system with two time scales, namely the van der Pol oscillator [12], which is a single parameter system

and hence not amenable to sloppy model analysis. Instead, by choosing appropriate perturbations  $\mathbf{g}(\mathbf{z})$ , we calculate the susceptibility of its dynamics: We make perturbations on the attractor and then systematically increase the separation of time scales in its dynamics to show how it can generally enhance the sloppiness in nonlinear systems.

## II. MULTIPLE-TIME-SCALE DYNAMICS

Multiple time scales are often found in the solutions of dynamical systems [13]. Broadly speaking, the defining criterion of these models is that the trajectory of one or more phase variables has an identifiable fast piece, such as a jump or a pulse, and a slow piece where the value of the variable does not change quickly [14]. In two dimensions, these systems are commonly studied in the context of slow-fast vector fields written as

$$\epsilon \dot{x} = X(x, y, \epsilon), \quad \dot{y} = Y(x, y, \epsilon), \quad (1)$$

where the parameter  $\epsilon > 0$  is small and the overdot indicates derivative with respect to time  $t$ . For  $O(1)$  functions  $X$  and  $Y$ , and  $X \neq 0$ :  $\dot{x} = O(1/\epsilon)$  and  $\dot{y} = O(1)$ , so that  $\epsilon$  is the ratio of time scales in the system. On one extreme, the singular limit  $\epsilon = 0$  corresponds to a differential algebraic system  $X = 0$ ,  $Y = \dot{y}$ , where the solutions of  $X = 0$  comprise the “critical manifold” close to which the flow in phase space is slow (the “slow manifold”). Similarly,  $\epsilon = 1$  corresponds to a limit where there is no separation of time scales, with a crossover at intermediate values of  $\epsilon$ .

Originally introduced in 1927, the van der Pol equation,  $\ddot{x} - \mu(1 - x^2)\dot{x} + x = 0$ , is a well-studied example of a second-order, nonlinear system with multiple time scales in its solution. Using the Liénard transformation  $y = x - x^3/3 - \dot{x}/\mu$ , and redefining time  $t \rightarrow t\mu$ , the equation can be written as a two-dimensional system [14,15] given by

$$\mu^{-2}\dot{x} = x - \frac{x^3}{3} - y, \quad \dot{y} = x, \quad (2)$$

which has the same form as (1) with  $\epsilon = \mu^{-2}$ . The global attractor of this dynamical system is a structurally stable limit cycle with two time scales [16].

\*sethna@lassp.cornell.edu




FIG. 1. (Color online) (a) Eigenvalues of the Hessian matrix of the cost of fitting at  $\mu = 1$ . (b) and (c) Top row: one period of time series  $x(\tau)$  (dotted line) and  $y(\tau)$  (solid line) for  $0 < \tau < 1$  are shown for  $\mu = 1$  and 100 as a function of time along with schematic error bars for the data-fitting of the trajectory of variable  $y$ . (d) Eigenvalues for  $\mu = 100$  are shown on the right. As  $\mu \rightarrow \infty$ , the orbit collapses onto the critical manifold with the trajectory spending most of its time on the slow manifold and vanishingly short on the jumps. Also shown in (b) and (c), bottom row, is the orbit in the  $xy$  plane (solid line) and the critical manifold (dashed line).

The van der Pol system provides a convenient way to separate time scales by varying  $\mu$ : Small values of  $\mu$  in the van der Pol system correspond to a small separation of time scales. It can be shown that the trajectory approaches that of the harmonic oscillator as  $\mu \rightarrow 0$  [12]. At large values of  $\mu$ , the system shows a separation of time scales which increases with increasing  $\mu$ . As shown in Figs. 1(b) and 1(c), with increasing  $\mu$ , the trajectory of  $x$  separates into a slow part that lies  $O(\mu^{-2})$  close to the phase space curve given by  $\dot{x} = 0$ , i.e., the critical manifold  $y = x - x^3/3$ , and a fast part which connects the two branches of the slow flow. Likewise, the separation of time scales in  $y$  is associated with the increasing sharpness of the kink in its trajectory.

The fact that with an increasing separation of time scales the trajectory spends an increasing amount time on the slow manifold and a decreasing amount of time on the jumps has important implications for fitting parameters to time series data of the van der Pol system. With increasing scale separation, one expects that the cost of fitting will be decreasingly sensitive to changes in the jumps of the trajectory as they get progressively shorter in duration.

### III. SLOPPINESS IN NONLINEAR FITS

In this section, we discuss the concepts of sloppiness and structural susceptibility in more detail with examples as a prelude to our calculations. For time series  $z(t, \mathbf{a})$ , a least-squares fit to data  $d_i$  minimizes a cost  $C = \frac{1}{2} \sum_i [z(t_i, \mathbf{a}) - d_i]^2 / \sigma_i^2$  in the space of system parameters, which are collectively denoted as  $\mathbf{a}$ . Our discovery of sloppiness is essentially that the eigenvalues of the Hessian of the cost with respect to

parameters,  $\mathcal{H}_{\alpha\beta} = \partial^2 C / \partial a_\alpha \partial a_\beta$ , at the best fit span many orders of magnitude. The larger and smaller eigenvalues correspond to stiffer and sloppier directions, respectively. For concreteness, consider a time series of a multiparameter model, such as the one denoted by  $y(\tau)$  in Fig. 1(b), top row. The error bars schematically show the least-squares fit of  $y(\tau)$ , and the side bar [Fig. 1(a)] shows the eigenvalues of the corresponding Hessian matrix. Note the broad range of eigenvalues ( $\sim 10^{11}$ , corresponding to a factor of almost a million in parameter range)—a feature that turns out to be typical in nonlinear fits.

Another vivid example of sloppiness in nonlinear models is provided by the well-established formalism behind the characterization of the sensitivities of initial conditions using Lyapunov exponents [17]. Consider  $dz/dt = \mathbf{f}(\mathbf{z})$  as a model whose parameters are the initial conditions  $a_\alpha = z_\alpha(0)$  and whose predictions are the final positions  $z_i(t)$  at time  $t$ . At the best fit,  $\mathcal{H}_{\alpha\beta} = (J^T J)_{\alpha\beta}$ , where  $J_{i\alpha} = \partial z_i(t) / \partial z_\alpha(0)$  is the Jacobian of the sensitivities to perturbations in the initial conditions. The Lyapunov exponents, which are defined to be the eigenvalues  $\ell_n$  of  $\mathbf{L} = \lim_{t \rightarrow \infty} 1/(2t) \log(J^T J)$ , utilize the same Hessian we would use in calculating the sloppy model eigenvalues  $\lambda_n = \exp(2t\ell_n)$ . The roughly equal spacing of Lyapunov exponents naturally explains both the exponentially broad range of sloppy model exponents and the roughly equal spacing of  $\log(\lambda_n)$  for a model with initial conditions as parameters.

Instead of the sensitivities with respect to the initial conditions or other intrinsic parameters, we focus here on the sensitivity of the dynamics to changes in the dynamical evolution laws. Therefore, for the remainder of this paper we will be interested in a cost function that measures the square of the distance between two time series for the system  $dz/dt = \mathbf{f}(\mathbf{z}) + \mathbf{a} \cdot \mathbf{g}(\mathbf{z})$ —one with perturbations, i.e.,  $\mathbf{z}(t, \mathbf{a} \rightarrow \mathbf{0})$ , and the other one without, i.e.,  $\mathbf{z}(t, \mathbf{a} = \mathbf{0})$ ,

$$C = \frac{1}{2} \int_0^T \|\mathbf{z}(t, \mathbf{a} \rightarrow \mathbf{0}) - \mathbf{z}(t, \mathbf{a} = \mathbf{0})\|^2 dt, \quad (3)$$

with the perturbing terms  $g_i(\mathbf{z})$  giving a power series in the components of  $\mathbf{z}$ . Later in the paper, we will use this form of the cost to compute the susceptibility of the van der Pol system and show how sloppiness is enhanced by increasing the separation of time scales in the van der Pol equations. This is in essence captured by Figs. 1(a) and 1(d), where we show that an increase in the van der Pol parameter  $\mu$  from 1 to 100 produces roughly a millionfold increase in the spread of eigenvalues.

### IV. SUSCEPTIBILITY OF THE van der POL SYSTEM

We perturb the van der Pol system in (2) by adding a series of additional terms. There is a long tradition in dynamical systems of studying equations of motion of polynomial form [17,18]; indeed, the theory of normal forms suggests that general dynamical systems, even at bifurcations, can be generically mapped into a polynomial form by a nonlinear but smooth change of variables. Adding extra polynomial terms is routinely done to “unfold” the qualitative behavior near bifurcations [19]. Here we focus on quantitative changes far from bifurcations. In choosing our perturbations, we must cut off the polynomials at some order. There are two ways in




FIG. 2. (Color online) Eigenvalues of the Hessian matrix are shown here as a function of  $\mu$ . The range  $1 \leq \mu \leq 100$  corresponds to a ratio of time scales  $1 \leq \epsilon \leq 10\,000$ . The five largest eigenvalues (solid lines) correspond to stiff directions in the parameter space: these directions perturb the slow manifold. The remainder (dashed lines) affect the transient part of the trajectory, which becomes smaller with an increasing separation of time scales, and hence these directions are decreasingly relevant.

which we specialize our general susceptibility analysis to the two-time-scale, periodic limit cycle of the van der Pol system. First, we choose the family of perturbations of order  $3N$  as follows:

$$\begin{aligned} \mu^{-2}\dot{x} &= x - \frac{x^3}{3} - y + \sum_{m+n \leq N} a_{m,n} \left( x - \frac{x^3}{3} - y \right)^m x^n, \\ \dot{y} &= x. \end{aligned} \quad (4)$$

This choice has two noteworthy features. (a) We have grouped the polynomial perturbations so that, for  $m \neq 0$ , they vanish on the critical manifold,  $y = x - x^3/3$ . That is, the parameters  $a_{m,n}$  with  $m \neq 0$  do not significantly affect the dynamics on the slow manifold; we call these the “fast parameters,” and correspondingly the  $a_{0,n}$  are “slow parameters.” The parameter  $a_{1,0}$  duplicates  $\mu$  to the same effect as changing the period, and we omit it. Surely, the eigenvalue spectrum of the general polynomial expansion,  $a_{m,n}x^m y^n$ , behaves qualitatively

similarly to the one presented here, but our parametrization greatly simplifies the analysis of the eigenvector perturbations. (b) We only perturb the  $\dot{x}$  equation. Our choice corresponds to a general expansion of a second-order equation, with the acceleration  $\ddot{y} = \dot{x}$  written as a polynomial in the position  $y$  and velocity  $\dot{y} = x$ . Perturbing both equations produces similar behavior.

Second, we modified the cost to focus on the limit cycle of the van der Pol system in two ways: (a) by rescaling all trajectories in our analysis so that they have the same unit period, and (b) by changing the initial condition so that it lies on the perturbed orbit and that the perturbed and unperturbed orbits are in phase with each other [20]. When we correct the period  $T$  by  $\delta T$ , initial conditions  $\mathbf{y}_0$  by  $\delta \mathbf{y}_0$ , and do an overall rescaling of the time variable  $t \rightarrow \tau T$ , the cost functional for the time series of  $y(\tau)$  at each  $\mu$  takes the following form:

$$\begin{aligned} C(\mu) &= \frac{1}{2} \int_0^1 [y(\tau, \mathbf{a} + \delta \mathbf{a}, \mathbf{y}_0 + \delta \mathbf{y}_0, T + \delta T) \\ &\quad - y(\tau, \mathbf{a}, \mathbf{y}_0, T)]^2 d\tau. \end{aligned} \quad (5)$$

In principle, changes in both time series,  $x(\tau)$  and  $y(\tau)$ , could be incorporated in the cost function, but we get qualitatively similar results by keeping either or both variables. Choosing to measure changes only in  $y(\tau)$  corresponds again to studying the second-order equation for  $\ddot{y}$  as an expansion in  $y$  and  $\dot{y}$ .

The susceptibilities are still given by the Hessian matrix at the best fit ( $\mathbf{a} = \mathbf{0}$ ):

$$\mathcal{H}(\mu)_{\alpha\beta} = \frac{\partial^2 C(\mu)}{\partial a_\alpha \partial a_\beta}, \quad (6)$$

which can be written out more completely as

$$\begin{aligned} \mathcal{H}(\mu)_{\alpha\beta} &= \int_0^1 \left( \frac{\partial y}{\partial a_\alpha} + \frac{\partial y}{\partial \mathbf{y}_0} \frac{\partial \mathbf{y}_0}{\partial a_\alpha} + \frac{\partial y}{\partial T} \frac{\partial T}{\partial a_\alpha} \right) \\ &\quad \times \left( \frac{\partial y}{\partial a_\beta} + \frac{\partial y}{\partial \mathbf{y}_0} \frac{\partial \mathbf{y}_0}{\partial a_\beta} + \frac{\partial y}{\partial T} \frac{\partial T}{\partial a_\beta} \right) d\tau. \end{aligned}$$




FIG. 3. (Color online) Hessian eigenvectors are shown for  $\mu = 1, 10$ , and  $100$ . Each colored small square shows the magnitude of an eigenvector component (the scale bar is shown on the right). Eigenvectors for each  $\mu$  are sorted so that the stiffer ones appear on the left; individual components are sorted so that “slow parameters” appear on the top. Note that with increasing  $\mu$ , the stiff and sloppy eigenvectors are separated by parameters: The stiff eigenvectors only have projections along the slow parameters, which perturb the slow manifold, whereas the sloppy directions have projections along the fast parameters, which mainly perturb the jumps.

Here, each of the two terms in the integral is to be interpreted as a Jacobian matrix, a mapping from the finite-dimensional parameter space to the infinite-dimensional data space:

$$J_{\tau\alpha} = \frac{\partial y(\tau)}{\partial a_\alpha} + \frac{\partial y(\tau)}{\partial \mathbf{y}_0} \frac{\partial \mathbf{y}_0}{\partial a_\alpha} + \frac{\partial y(\tau)}{\partial T} \frac{\partial T}{\partial a_\alpha}. \quad (7)$$

The sensitivity trajectories in the Jacobian— $\partial y/\partial a_\alpha$ ,  $\partial y/\partial \mathbf{y}_0$ , and  $\partial y/\partial T$ —were computed using the open source SLOPPY-CELL package [21,22]. The expressions for the time invariant quantities,  $\partial \mathbf{y}_0/\partial a_\alpha$  and  $\partial T/\partial a_\alpha$ , were found by enforcing periodicity of the perturbed time series denoted by  $\mathbf{y}(\tau) \equiv (x(\tau), y(\tau))$  as follows:

$$\begin{aligned} \mathbf{y}(\tau = 0, \mathbf{a} + \delta \mathbf{a}, \mathbf{y}_0 + \delta \mathbf{y}_0, T + \delta T) \\ = \mathbf{y}(\tau = 1, \mathbf{a} + \delta \mathbf{a}, \mathbf{y}_0 + \delta \mathbf{y}_0, T + \delta T). \end{aligned}$$

Taylor expansion of both sides of the previous equation leads to a vector equation:

$$\delta \mathbf{y}_0 = \frac{\partial \mathbf{y}}{\partial T} \Big|_{\tau=1} \delta T + \frac{\partial \mathbf{y}}{\partial \mathbf{a}} \Big|_{\tau=1} \delta \mathbf{a} + \frac{\partial \mathbf{y}}{\partial \mathbf{y}_0} \Big|_{\tau=1} \delta \mathbf{y}_0,$$

from which both constants can be computed following the convention that the component denoting the change in the initial condition of  $y(\tau)$  in  $\delta \mathbf{y}_0$  is set to zero. Now with the Jacobian calculated, the Hessian at best fit is simply  $\mathcal{H} = J^T J$ .

### A. Eigenvalues and eigenvectors

We computed the Hessian matrix given by the previous equation at multiple values of  $\mu$ . The spread of eigenvalues (Fig. 2) increases as a function of  $\mu$  confirming that sloppiness increases with an increasing separation of time scales. Not surprisingly [23], for  $N = 4$ , the 14 eigenvalues for  $\mu = 1$  already span 11 orders of magnitude, while for  $\mu = 100$ , we observe that the stiffest eigenvalue is 18 orders of magnitude larger than the smallest one—the spread increases by  $10^7$  when  $\mu$  increases to 100.

Taken together with the eigenvectors shown in Fig. 3, some interesting facts come to light: Figure 2 shows that with increasing  $\mu$ , the eigenvalues separate into two clusters of closely related decay exponents. The largest  $N$  eigenvalues




FIG. 4. (Color online) Top three rows: Eigenpredictions  $\delta y_k$  for  $k = 0, 3, 6, 9, 12$  at  $\mu = 1, 10$ , and  $100$  are shown as solid red lines for stiff modes and dashed green lines for sloppy modes. These curves show the response of perturbations if the parameters are changed infinitesimally along the Hessian eigenvectors: A parameter change of norm  $\epsilon$  along eigendirection  $n$  will change the trajectories by  $\lambda_n \epsilon$  times the eigenprediction. Dotted gray lines show the unperturbed van der Pol solution for comparison ( $y$  scale on the right-hand side). As the time scales separate, the amplitudes of the sloppiest eigenpredictions increase (roughly in proportion to  $\mu$ ), becoming increasingly concentrated at the jumps. The bottom row shows the eigencycles for  $\mu = 100$  as solid red lines and green dashed lines corresponding to the perturbations in row 3 (i.e., the new limit cycle for a perturbation of strength  $\epsilon \sim 1/\lambda_n$ ). These curves show how the van der Pol orbit changes with perturbations along the Hessian eigenvectors. Both the stiff and the sloppy modes change the orbit at the jumps (occurring at the extrema in the dashed lines); the stiff modes also change behavior at the slow manifold, whereas the sloppy modes only affect the jumps.

approach constants. The other eigenvalues decay with power laws: two modes with exponents between  $-2$  and  $-3$  and the remaining modes with exponents between  $-5$  and  $-6$ . Similarly, Fig. 3 shows that the eigenvectors also separate into two groups with increasing  $\mu$ : The stiffest directions are linear combinations of the slow parameters, whereas the sloppy directions are comprised of other parameters, as expected.

We can understand the effect of perturbations in parameter combinations given by the eigenvectors (called *eigenparameters*)  $\hat{e}_k$  more clearly by observing their behavior in the data space. The Jacobian transformation of (7) projects the eigenvectors to the data space:  $\delta y_k = J \cdot \hat{e}_k / \sqrt{\lambda_k}$ , where  $\lambda_k$  corresponds to the  $k$ th largest eigenvalue. Defined in this way, these data space vectors, called *eigenpredictions* [4],  $\delta y_k$ , are also orthonormal. Alternatively, the eigenpredictions are the left singular vectors in the singular value decomposition of the Jacobian (i.e., they are the columns of the unitary matrix  $U$  in  $J = U \Sigma V^T$  [24]). As shown in Fig. 4 for  $\mu = 1, 10$ , and  $100$  (top three rows), we learn from the eigenpredictions that the stiff modes affect behavior both along the slow manifold and at the jumps. Moreover, with increasing  $\mu$ , as the eigenvalues associated with the stiff directions approach constants (Fig. 2), so do the stiff eigenpredictions [Fig. 4, rows 2 and 3, columns (a) and (b)]. The sloppy modes, on the other hand, affect the dynamics on the jumps only. The maximum amplitudes of the (normalized) sloppiest eigenpredictions appear to increase roughly proportional to  $\mu$  (corresponding to the jump duration of  $\sim \mu^{-2}$ ). In the limit, these become  $\delta$  functions and derivatives concentrated at the jumps. Figure 4 (bottom row) also shows the limit cycles (*eigencycles*) with eigenparameter perturbations as phase space trajectories  $(x, y + \eta \delta y_k)$  for small  $\eta$ .

## V. DISCUSSION

In this paper, we have introduced a formalism that we call “structural susceptibility” for analyzing the quantitative dependence of dynamical systems to perturbations of the equations of motion. It is a generalization of “unfolding” methods of bifurcation theory and the Lyapunov exponents governing the dependence on initial conditions, and it exposes the ubiquitous presence of a broad range of sloppy eigendirections in parameter space—largely unimportant to the dynamics. We used this method to study the role of time-scale separation in enhancing the sloppiness of the susceptibility spectrum in the particular case of the van der Pol oscillator.

By extending the framework of our sloppy model analysis to systems where changes in evolution laws are to be studied, our method offers a simple way to calculate the effects of broad classes of perturbations. By studying the structural susceptibility of a dynamical system with two time scales, the analysis presented here showed that the sloppiness of nonlinear

systems is enhanced by a separation of time scales in the dynamics. With increasing separation of time scales in the van der Pol oscillator, the trajectory spends an increasing amount of time on the slow manifold and a vanishingly small amount of time in the transition region. The cost of perturbations is integrated over time and therefore we are unsurprised that the perturbations that change the slow manifold will accrue the most cost and therefore manifest as stiff modes of the Hessian matrix. The remaining directions are sloppy as they only affect the behavior at the jumps or the fast pieces. These perturbations manifest as  $\delta$  functions and their derivatives—significantly affecting the phase-space trajectory, but over only the fast times asymptotically ignored in the least-squares cost. It remains a challenge to connect the separation of time scales to parameter sensitivity in more complicated systems, but the analogy of the van der Pol system’s behavior with other nonlinear physical systems of interest is clear.

Many important dynamical systems have multiple time scales in their solutions: examples include models in neuroscience (such as Hodgkin-Huxley model), systems biology or chemical reaction systems (such as protein network models), and engineering (such as models of combustion, lasers, locomotion, etc.). Our analysis suggests that any system with multiple time scales should become sloppier as the scales separate for the same reasons that we found in the van der Pol system: Some parameter combinations will only affect the fast dynamics and lead to sloppy modes. Perturbations which affect the slow dynamics will presumably accrue more cost and be stiff.

More broadly, the sloppiness exposed by our structural susceptibility analysis has clear implications for attempting to reconstruct the equations of motion from experimental data [25] because the true dynamics along any sloppy eigendirection will be relatively poorly determined. This discovery has already influenced work on experimental design optimization: estimating parameters is challenging [8,26], but extracting predictions without constraining parameters is straightforward [6]. We further anticipate that the concept of structural susceptibility will be useful for studying systems with chaos, bifurcations, and phase transitions; quantifying the unfoldings of these systems may also be useful for gaining a deeper understanding of the phenomena they model.

## ACKNOWLEDGMENTS

We thank Stefanos Papanikolaou for helpful conversations that led us to think about perturbing the slow manifold and about the analogs to thermodynamic susceptibilities, and John Guckenheimer for valuable insights and discussions regarding our calculations. Support from NSF Grant No. DMR 1005479 is gratefully acknowledged.

[1] We employ the word *structural* in the same context as its usage in dynamical systems literature on *structural stability*. The word *susceptibility* is inspired from physics wherein it is a measure of response to a perturbation (such as an applied external field) quantified by the second derivative of the free

energy with respect to parameters. Since cost is analogous to free energy (in that both are minimized), it is natural to refer to the response to perturbations in dynamics, also quantified via second derivatives, as *structural susceptibility*.

[2] K. S. Brown and J. P. Sethna, *Phys. Rev. E* **68**, 021904 (2003).

- [3] R. N. Gutenkunst, J. J. Waterfall, F. P. Casey, K. S. Brown, C. R. Myers, and J. P. Sethna, *PLoS Comput Biol* **3**, e189 (2007).
- [4] M. K. Transtrum, B. B. Machta, and J. P. Sethna, *Phys. Rev. E* **83**, 036701 (2011).
- [5] F. P. Casey, D. Baird, Q. Feng, R. N. Gutenkunst, J. J. Waterfall, C. R. Myers, K. S. Brown, R. A. Cerione, and J. P. Sethna, *IET Syst. Biol.* **1**, 190 (2007).
- [6] R. N. Gutenkunst, F. P. Casey, J. J. Waterfall, C. R. Myers, and J. P. Sethna, *Ann. N.Y. Acad. Sci.* **1115**, 203 (2007).
- [7] M. K. Transtrum, B. B. Machta, and J. P. Sethna, *Phys. Rev. Lett.* **104**, 060201 (2010).
- [8] J. F. Apgar, D. K. Witmer, F. M. White, and B. Tidor, *Mol. BioSyst.* **6**, 1890 (2010).
- [9] M. Secrier, T. Toni, and M. P. H. Stumpf, *Mol. BioSyst.* **5**, 1925 (2009).
- [10] A. Dayarian, M. Chaves, E. D. Sontag, and A. M. Sengupta, *PLoS Comput Biol* **5**, e1000256 (2009).
- [11] H. Hettling and J. H. G. M. van Beek, *PLoS Comput Biol* **7**, e1002130 (2011).
- [12] B. van der Pol, *The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science Ser. 7*(2), 978 (1926).
- [13] *Multiple-time-scale Dynamical Systems*, edited by C. K. R. T. Jones and A. I. Khibnik (Springer, New York, NY, 2000).
- [14] J. Grasman, *Asymptotic Methods for Relaxation Oscillations and Applications* (Springer, New York, NY, 1987).
- [15] S. H. Strogatz, *Nonlinear Dynamics and Chaos: With Applications to Physics, Biology, Chemistry and Engineering* (Westview, Boulder, CO, 2001).
- [16] Incidentally, the set of equations (2) can also be considered to be a special case of the FitzHugh-Nagumo model [see R. FitzHugh, *Biophys J.* **1**, 445 (1961)] introduced three decades later as a simplification of the Hodgkin-Huxley equations of neuronal spikes in the squid giant axons, and is sometimes referred to as the Bonhoeffer–van der Pol model.
- [17] A. Katok and B. Hasselblatt, *Introduction to the Modern Theory of Dynamical Systems* (Cambridge University Press, New York, NY, 1997).
- [18] J. M. Guckenheimer and P. Holmes, *Nonlinear Oscillations, Dynamical Systems and Bifurcations of Vector Fields* (Springer, New York, NY, 1983).
- [19] J. Murdock, *Normal Forms and Unfoldings for Local Dynamical Systems* (Springer, New York, 2003).
- [20] Perturbations distort the dynamics so that the attractor and its period change. We addressed these issues by setting the periods to unity, and by moving the initial conditions to the new attractor to remove any transients. Alternatively, if we fit data over many periods without making the said changes, the parameter combinations determining the period and phase would become stiff modes in our dynamics.
- [21] R. N. Gutenkunst, J. C. Atlas, F. P. Casey, R. S. Kuczynski, J. J. Waterfall, C. R. Myers, and J. P. Sethna, SLOPPYCELL, <http://sloppycell.sourceforge.net> (2007).
- [22] C. R. Myers, R. N. Gutenkunst, and J. P. Sethna, *Comput. Sci. Eng.* **9**, 34 (2007).
- [23] We understand this as sloppiness as arising due to the generalized interpolation argument [4].
- [24] W. H. Press, S. A. Teukolsky, W. T. Vetterling, and B. P. Flannery, *Numerical Recipes: The Art of Scientific Computing*, 3rd ed. (Cambridge University Press, New York, 2007).
- [25] J. Bongard and H. Lipson, *Proc. Natl. Acad. Sci. (USA)* **104**, 9943 (2007).
- [26] R. Chachra, M. K. Transtrum, and J. P. Sethna, *Mol. BioSyst.* **7**, 2522 (2011).