

Parallel Equations for the Electric and Magnetic Fields

Dr. Colton, Physics 441, Fall 2016

ELECTRIC

Statics

$$1. \quad q = \int \lambda dl$$
$$q = \int \sigma da$$
$$q = \int \rho d\tau$$

$$2. \quad \mathbf{F} = \frac{1}{4\pi\epsilon_0} \frac{qQ}{|\mathbf{r}-\mathbf{r}'|^3} (\mathbf{r} - \mathbf{r}')$$

$$3. \quad \mathbf{F} = Q\mathbf{E}$$

$$4. \quad \mathbf{E}(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \int \frac{dq}{|\mathbf{r}-\mathbf{r}'|^3} (\mathbf{r} - \mathbf{r}')$$

$$\mathbf{E}(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \int \frac{\lambda(\mathbf{r}')d\mathbf{l}'}{|\mathbf{r}-\mathbf{r}'|^3} (\mathbf{r} - \mathbf{r}')$$

$$\mathbf{E}(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \int_S \frac{\sigma(\mathbf{r}')d\mathbf{a}'}{|\mathbf{r}-\mathbf{r}'|^3} (\mathbf{r} - \mathbf{r}')$$

$$\mathbf{E}(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \int_V \frac{\rho(\mathbf{r}')d\tau'}{|\mathbf{r}-\mathbf{r}'|^3} (\mathbf{r} - \mathbf{r}')$$

$$5. \quad \nabla \cdot \mathbf{E} = \frac{\rho}{\epsilon_0}$$

$$6. \quad \Phi_E = \int_S \mathbf{E} \cdot d\mathbf{a}$$

$$7. \quad \oint_S \mathbf{E} \cdot d\mathbf{a} = \frac{Q_{enc}}{\epsilon_0}$$

$$8. \quad \nabla \times \mathbf{E} = 0 \quad (\text{this gets modified below})$$

$$9. \quad \mathbf{E} = -\nabla V \quad (\text{this gets modified in Phys 442})$$

$$10. \quad V(\mathbf{r}) = -\int_{ref}^{\mathbf{r}} \mathbf{E} \cdot d\mathbf{l}$$

$$11. \quad V(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \int \frac{\lambda(\mathbf{r}')}{|\mathbf{r}-\mathbf{r}'|} dl'$$

$$V(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \int \frac{\sigma(\mathbf{r}')}{|\mathbf{r}-\mathbf{r}'|} da'$$

$$V(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \int \frac{\rho(\mathbf{r}')}{|\mathbf{r}-\mathbf{r}'|} d\tau'$$

$$12. \quad U = \frac{\epsilon_0}{2} \int E^2 d\tau$$

$$13. \quad C = \frac{Q}{V}$$

$$14. \quad U = \frac{1}{2} \frac{Q^2}{C}$$

$$15. \quad E_1^\perp - E_2^\perp = \frac{\sigma}{\epsilon_0}$$

$$16. \quad \mathbf{E}_1^\parallel = \mathbf{E}_2^\parallel$$

$$17. \quad V_1 = V_2$$

$$18. \quad \nabla^2 V = -\frac{\rho}{\epsilon_0}$$

MAGNETIC

Statics

$$1. \quad I = \int K_\perp dl$$
$$I = \int \mathbf{J} \cdot d\mathbf{a}$$

$$2. \quad \text{No easy parallel for magnetic field}$$

$$3. \quad \mathbf{F} = Q(\mathbf{v} \times \mathbf{B})$$

$$4. \quad \mathbf{B}(\mathbf{r}) = \frac{\mu_0}{4\pi} \int \frac{\mathbf{l} \times (\mathbf{r}-\mathbf{r}')}{|\mathbf{r}-\mathbf{r}'|^3} dl'$$

$$\mathbf{B}(\mathbf{r}) = \frac{\mu_0}{4\pi} \int_S \frac{\mathbf{K}(\mathbf{r}') \times (\mathbf{r}-\mathbf{r}')}{|\mathbf{r}-\mathbf{r}'|^3} da'$$

$$\mathbf{B}(\mathbf{r}) = \frac{\mu_0}{4\pi} \int_V \frac{\mathbf{J}(\mathbf{r}') \times (\mathbf{r}-\mathbf{r}')}{|\mathbf{r}-\mathbf{r}'|^3} d\tau'$$

$$5. \quad \nabla \cdot \mathbf{B} = 0$$

$$6. \quad \Phi_B = \int_S \mathbf{B} \cdot d\mathbf{a}$$

$$7. \quad \oint \mathbf{B} \cdot d\mathbf{l} = \mu_0 I_{enc}$$

$$8. \quad \nabla \times \mathbf{B} = \mu_0 \mathbf{J} \quad (\text{this gets modified below})$$

$$9. \quad \mathbf{B} = \nabla \times \mathbf{A}, \quad \nabla \cdot \mathbf{A} = 0 \quad (\text{Coulomb gauge})$$

$$10. \quad \text{No direct parallel for the magnetic field}$$

$$11. \quad \mathbf{A}(\mathbf{r}) = \frac{\mu_0}{4\pi} \int \frac{\mathbf{l}}{|\mathbf{r}-\mathbf{r}'|} dl'$$

$$\mathbf{A}(\mathbf{r}) = \frac{\mu_0}{4\pi} \int \frac{\mathbf{K}}{|\mathbf{r}-\mathbf{r}'|} da'$$

$$\mathbf{A}(\mathbf{r}) = \frac{\mu_0}{4\pi} \int \frac{\mathbf{J}(\mathbf{r}')}{|\mathbf{r}-\mathbf{r}'|} d\tau'$$

$$12. \quad U = \frac{1}{2\mu_0} \int B^2 d\tau$$

$$13. \quad L = \frac{\Phi}{I}$$

$$14. \quad U = \frac{1}{2} LI^2$$

$$15. \quad B_1^\perp = B_2^\perp$$

$$16. \quad \mathbf{B}_1^\parallel - \mathbf{B}_2^\parallel = \mu_0 \mathbf{K}$$

$$17. \quad \mathbf{A}_1 = \mathbf{A}_2$$

$$18. \quad \nabla^2 \mathbf{A} = -\mu_0 \mathbf{J}$$

$$19. V(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \sum_{n=0}^{\infty} \frac{1}{r^{n+1}} \int (r')^n P_n(\cos\theta') \rho(\mathbf{r}') d\tau'$$

Materials

$$20. \mathbf{p} = \int \mathbf{r}' \rho(\mathbf{r}') d\tau'$$

$$21. V_{\text{dip}} = \frac{1}{4\pi\epsilon_0} \frac{\mathbf{p} \cdot \hat{\mathbf{r}}}{r^2}$$

$$22. \mathbf{E}_{\text{dip}}(r, \theta) = \frac{1}{4\pi\epsilon_0} \frac{p}{r^3} (2\cos\theta \hat{\mathbf{r}} + \sin\theta \hat{\boldsymbol{\theta}})$$

$$23. \mathbf{N} = \mathbf{p} \times \mathbf{E}$$

$$24. \mathbf{F} = (\mathbf{p} \cdot \nabla) \mathbf{E}$$

$$25. U = -\mathbf{p} \cdot \mathbf{E}$$

$$26. \mathbf{P} = \text{dipole moment per unit volume}$$

$$27. \mathbf{D} = \epsilon_0 \mathbf{E} + \mathbf{P}$$

$$28. \nabla \cdot \mathbf{D} = \rho_f$$

$$29. \mathbf{P} = \epsilon_0 \chi_e \mathbf{E}$$

$$30. \mathbf{D} = \epsilon \mathbf{E} = \epsilon_0 \epsilon_r \mathbf{E}$$

$$31. \sigma_b = \mathbf{P} \cdot \hat{\mathbf{n}}$$

$$32. \rho_b = -\nabla \cdot \mathbf{P}$$

$$33. \oint \mathbf{D} \cdot d\mathbf{a} = Q_{f,enc}$$

$$34. \epsilon_r = 1 + \chi_e = \frac{\epsilon}{\epsilon_0}$$

$$35. U = \frac{1}{2} \int \mathbf{D} \cdot \mathbf{E} d\tau$$

Dynamics

$$37. \nabla \cdot \mathbf{J} = -\frac{\partial \rho}{\partial t}$$

$$38. \nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t}$$

$$39. \nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t} \quad (\text{unchanged for materials})$$

$$19. \mathbf{A}(\mathbf{r}) = \frac{\mu_0 I}{4\pi} \sum_{n=0}^{\infty} \frac{1}{r^{n+1}} \oint (r')^n P_n(\cos\theta') d\mathbf{l}'$$

Materials

$$20. \mathbf{m} = I \int d\mathbf{a} = I \mathbf{a}$$

$$21. \mathbf{A}_{\text{dip}} = \frac{\mu_0}{4\pi} \frac{\mathbf{m} \times \hat{\mathbf{r}}}{r^2}$$

$$22. \mathbf{B}_{\text{dip}}(r, \theta) = \frac{\mu_0 m}{4\pi r^3} (2\cos\theta \hat{\mathbf{r}} + \sin\theta \hat{\boldsymbol{\theta}})$$

$$23. \mathbf{N} = \mathbf{m} \times \mathbf{B}$$

$$24. \mathbf{F} = \nabla(\mathbf{m} \cdot \mathbf{B})$$

$$25. U = -\mathbf{m} \cdot \mathbf{B}$$

$$26. \mathbf{M} = \text{magnetic dipole moment per unit volume}$$

$$27. \mathbf{H} = \mathbf{B}/\mu_0 - \mathbf{M}$$

$$28. \nabla \cdot \mathbf{B} = 0 \quad (\text{still})$$

$$29. \mathbf{M} = \chi_m \mathbf{H}$$

$$30. \mathbf{H} = \mathbf{B}/\mu = \mathbf{B}/(\mu_0 \mu_r)$$

$$31. \mathbf{J}_b = \nabla \times \mathbf{M}$$

$$32. \mathbf{K}_b = \mathbf{M} \times \hat{\mathbf{n}}$$

$$33. \oint \mathbf{H} \cdot d\mathbf{l} = I_{f,enc}$$

$$34. \mu_r = 1 + \chi_m = \frac{\mu}{\mu_0}$$

$$35. U = \frac{1}{2} \int \mathbf{H} \cdot \mathbf{B} d\tau$$

Dynamics

$$37. \nabla \cdot \mathbf{J} = -\frac{\partial \rho}{\partial t} \quad (\text{same equation as in left hand column; connects charge to current})$$

$$38. \nabla \times \mathbf{B} = \mu_0 \mathbf{J} + \mu_0 \epsilon_0 \frac{\partial \mathbf{E}}{\partial t}$$

$$39. \nabla \times \mathbf{H} = \mu_0 \mathbf{J}_f + \frac{\partial \mathbf{D}}{\partial t}$$