

## Parallel Equations for the Electric and Magnetic Fields

Dr. Colton, Physics 441, Fall 2017

### ELECTRIC

#### Statics

1.  $q = \int \lambda dl$
- $q = \int \sigma da$
- $q = \int \rho d\tau$
2.  $\mathbf{F} = \frac{1}{4\pi\epsilon_0} \frac{qQ}{|\mathbf{r}-\mathbf{r}'|^3} (\mathbf{r} - \mathbf{r}')$
3.  $\mathbf{F} = Q\mathbf{E}$
4.  $\mathbf{E}(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \int \frac{dq}{|\mathbf{r}-\mathbf{r}'|^3} (\mathbf{r} - \mathbf{r}')$   

$$\mathbf{E}(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \int \frac{\lambda(\mathbf{r}')dl'}{|\mathbf{r}-\mathbf{r}'|^3} (\mathbf{r} - \mathbf{r}')$$
  

$$\mathbf{E}(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \int_S \frac{\sigma(\mathbf{r}')da'}{|\mathbf{r}-\mathbf{r}'|^3} (\mathbf{r} - \mathbf{r}')$$
  

$$\mathbf{E}(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \int_V \frac{\rho(\mathbf{r}')d\tau'}{|\mathbf{r}-\mathbf{r}'|^3} (\mathbf{r} - \mathbf{r}')$$

$$5. \boxed{\nabla \cdot \mathbf{E} = \frac{\rho}{\epsilon_0}}$$

6.  $\Phi_E = \int_S \mathbf{E} \cdot d\mathbf{a}$
7.  $\oint_S \mathbf{E} \cdot d\mathbf{a} = \frac{Q_{enc}}{\epsilon_0}$
8.  $\nabla \times \mathbf{E} = 0$  (this gets modified below)
9.  $\mathbf{E} = -\nabla V$  (this gets modified in Phys 442)
10.  $V(\mathbf{r}) = - \int_{\text{ref}}^{\mathbf{r}} \mathbf{E} \cdot d\mathbf{l}$
11.  $V(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \int \frac{\lambda(\mathbf{r}')}{|\mathbf{r}-\mathbf{r}'|} dl'$   

$$V(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \int \frac{\sigma(\mathbf{r}')}{|\mathbf{r}-\mathbf{r}'|} da'$$
  

$$V(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \int \frac{\rho(\mathbf{r}')}{|\mathbf{r}-\mathbf{r}'|} d\tau'$$

12.  $U = \frac{\epsilon_0}{2} \int E^2 d\tau$
13.  $C = \frac{Q}{V}$
14.  $U = \frac{1}{2} \frac{Q^2}{C}$
15.  $E_1^\perp - E_2^\perp = \frac{\sigma}{\epsilon_0}$
16.  $\mathbf{E}_1^\parallel = \mathbf{E}_2^\parallel$
17.  $V_1 = V_2$
18.  $\nabla^2 V = -\frac{\rho}{\epsilon_0}$

### MAGNETIC

#### Statics

1.  $I = \int K_\perp dl$   
 $I = \int \mathbf{J} \cdot d\mathbf{a}$
2. No easy parallel for magnetic field
3.  $\mathbf{F} = Q(\mathbf{v} \times \mathbf{B})$
4.  $\mathbf{B}(\mathbf{r}) = \frac{\mu_0}{4\pi} \int \frac{\mathbf{I} \times (\mathbf{r}-\mathbf{r}')}{|\mathbf{r}-\mathbf{r}'|^3} dl'$   

$$\mathbf{B}(\mathbf{r}) = \frac{\mu_0}{4\pi} \int_S \frac{\mathbf{K}(\mathbf{r}') \times (\mathbf{r}-\mathbf{r}')}{|\mathbf{r}-\mathbf{r}'|^3} da'$$
  

$$\mathbf{B}(\mathbf{r}) = \frac{\mu_0}{4\pi} \int_V \frac{\mathbf{J}(\mathbf{r}') \times (\mathbf{r}-\mathbf{r}')}{|\mathbf{r}-\mathbf{r}'|^3} d\tau'$$

$$5. \boxed{\nabla \cdot \mathbf{B} = 0}$$

6.  $\Phi_B = \int_S \mathbf{B} \cdot d\mathbf{a}$
7.  $\oint \mathbf{B} \cdot d\mathbf{l} = \mu_0 I_{enc}$
8.  $\nabla \times \mathbf{B} = \mu_0 \mathbf{J}$  (this gets modified below)
9.  $\mathbf{B} = \nabla \times \mathbf{A}, \nabla \cdot \mathbf{A} = 0$  (Coulomb gauge)
10. No direct parallel for the magnetic field
11.  $\mathbf{A}(\mathbf{r}) = \frac{\mu_0}{4\pi} \int \frac{\mathbf{I}}{|\mathbf{r}-\mathbf{r}'|} dl'$   

$$\mathbf{A}(\mathbf{r}) = \frac{\mu_0}{4\pi} \int \frac{\mathbf{K}}{|\mathbf{r}-\mathbf{r}'|} da'$$
  

$$\mathbf{A}(\mathbf{r}) = \frac{\mu_0}{4\pi} \int \frac{\mathbf{J}(\mathbf{r}')}{|\mathbf{r}-\mathbf{r}'|} d\tau'$$

12.  $U = \frac{1}{2\mu_0} \int B^2 d\tau$
13.  $L = \frac{\Phi}{I}$
14.  $U = \frac{1}{2} LI^2$
15.  $B_1^\perp = B_2^\perp$
16.  $\mathbf{B}_1^\parallel - \mathbf{B}_2^\parallel = \mu_0 \mathbf{K}$
17.  $\mathbf{A}_1 = \mathbf{A}_2$
18.  $\nabla^2 \mathbf{A} = -\mu_0 \mathbf{J}$

$$19. V(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \sum_{n=0}^{\infty} \frac{1}{r^{n+1}} \int (r')^n P_n(\cos\theta') \rho(\mathbf{r}') d\tau'$$

### Materials

$$20. \mathbf{p} = \int \mathbf{r}' \rho(\mathbf{r}') d\tau'$$

$$21. V_{\text{dip}} = \frac{1}{4\pi\epsilon_0} \frac{\mathbf{p} \cdot \hat{\mathbf{r}}}{r^2}$$

$$22. \mathbf{E}_{\text{dip}}(r, \theta) = \frac{1}{4\pi\epsilon_0} \frac{p}{r^3} (2\cos\theta \hat{\mathbf{r}} + \sin\theta \hat{\mathbf{\theta}})$$

$$23. \mathbf{N} = \mathbf{p} \times \mathbf{E}$$

$$24. \mathbf{F} = (\mathbf{p} \cdot \nabla) \mathbf{E}$$

$$25. U = -\mathbf{p} \cdot \mathbf{E}$$

26.  $\mathbf{P}$  = dipole moment per unit volume

$$27. \mathbf{D} = \epsilon_0 \mathbf{E} + \mathbf{P}$$

$$28. \boxed{\nabla \cdot \mathbf{D} = \rho_f}$$

$$29. \mathbf{P} = \epsilon_0 \chi_e \mathbf{E}$$

$$30. \mathbf{D} = \epsilon \mathbf{E} = \epsilon_0 \epsilon_r \mathbf{E}$$

$$31. \sigma_b = \mathbf{P} \cdot \hat{\mathbf{n}}$$

$$32. \rho_b = -\nabla \cdot \mathbf{P}$$

$$33. \oint \mathbf{D} \cdot d\mathbf{a} = Q_{f,enc}$$

$$34. \epsilon_r = 1 + \chi_e = \frac{\epsilon}{\epsilon_0}$$

$$35. U = \frac{1}{2} \int \mathbf{D} \cdot \mathbf{E} d\tau$$

### Dynamics

$$37. \nabla \cdot \mathbf{J} = -\frac{\partial \rho}{\partial t}$$

$$38. \boxed{\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t}}$$

$$39. \boxed{\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t}} \quad (\text{unchanged for materials})$$

$$19. \mathbf{A}(\mathbf{r}) = \frac{\mu_0 I}{4\pi} \sum_{n=0}^{\infty} \frac{1}{r^{n+1}} \phi(r')^n P_n(\cos\theta') d\mathbf{l}'$$

### Materials

$$20. \mathbf{m} = I \int d\mathbf{a} = I \mathbf{a}$$

$$21. \mathbf{A}_{\text{dip}} = \frac{\mu_0}{4\pi} \frac{\mathbf{m} \times \hat{\mathbf{r}}}{r^2}$$

$$22. \mathbf{B}_{\text{dip}}(r, \theta) = \frac{\mu_0 m}{4\pi r^3} (2\cos\theta \hat{\mathbf{r}} + \sin\theta \hat{\mathbf{\theta}})$$

$$23. \mathbf{N} = \mathbf{m} \times \mathbf{B}$$

$$24. \mathbf{F} = \nabla(\mathbf{m} \cdot \mathbf{B})$$

$$25. U = -\mathbf{m} \cdot \mathbf{B}$$

26.  $\mathbf{M}$  = magnetic dipole moment per unit volume

$$27. \mathbf{H} = \mathbf{B}/\mu_0 - \mathbf{M}$$

$$28. \boxed{\nabla \cdot \mathbf{B} = 0} \quad (\text{still})$$

$$29. \mathbf{M} = \chi_m \mathbf{H}$$

$$30. \mathbf{H} = \mathbf{B}/\mu = \mathbf{B}/(\mu_0 \mu_r)$$

$$31. \mathbf{J}_b = \nabla \times \mathbf{M}$$

$$32. \mathbf{K}_b = \mathbf{M} \times \hat{\mathbf{n}}$$

$$33. \oint \mathbf{H} \cdot d\mathbf{l} = I_{f,enc}$$

$$34. \mu_r = 1 + \chi_m = \frac{\mu}{\mu_0}$$

$$35. U = \frac{1}{2} \int \mathbf{H} \cdot \mathbf{B} d\tau$$

### Dynamics

$$37. \nabla \cdot \mathbf{J} = -\frac{\partial \rho}{\partial t} \quad (\text{same equation as in left hand column; connects charge to current})$$

$$38. \boxed{\nabla \times \mathbf{B} = \mu_0 \mathbf{J} + \mu_0 \epsilon_0 \frac{\partial \mathbf{E}}{\partial t}}$$

$$39. \boxed{\nabla \times \mathbf{H} = \mathbf{J}_f + \frac{\partial \mathbf{D}}{\partial t}}$$