

No time limit. Student calculators are allowed. One page of handwritten notes allowed (front & back). Books not allowed.

Name \_\_\_\_\_

*Instructions:* Please label & circle/box your answers. Show your work, where appropriate! And remember: **in any problems involving Gauss's Law, you should explicitly show your Gaussian surface.** For all problems, unless otherwise specified you may assume that you are dealing with **electrostatics**, i.e. the charges are not moving and the fields have come to equilibrium.

*Griffiths front and back covers*

VECTOR DERIVATIVES	VECTOR IDENTITIES
<p><b>Cartesian.</b> <math>d\mathbf{l} = dx \hat{\mathbf{x}} + dy \hat{\mathbf{y}} + dz \hat{\mathbf{z}}</math>; <math>d\tau = dx dy dz</math></p> <p><b>Gradient:</b> <math>\nabla f = \frac{\partial f}{\partial x} \hat{\mathbf{x}} + \frac{\partial f}{\partial y} \hat{\mathbf{y}} + \frac{\partial f}{\partial z} \hat{\mathbf{z}}</math></p> <p><b>Divergence:</b> <math>\nabla \cdot \mathbf{v} = \frac{\partial v_x}{\partial x} + \frac{\partial v_y}{\partial y} + \frac{\partial v_z}{\partial z}</math></p> <p><b>Curl:</b> <math>\nabla \times \mathbf{v} = \left( \frac{\partial v_z}{\partial y} - \frac{\partial v_y}{\partial z} \right) \hat{\mathbf{x}} + \left( \frac{\partial v_x}{\partial z} - \frac{\partial v_z}{\partial x} \right) \hat{\mathbf{y}} + \left( \frac{\partial v_y}{\partial x} - \frac{\partial v_x}{\partial y} \right) \hat{\mathbf{z}}</math></p> <p><b>Laplacian:</b> <math>\nabla^2 f = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2}</math></p> <p><b>Spherical.</b> <math>d\mathbf{l} = dr \hat{\mathbf{r}} + r d\theta \hat{\boldsymbol{\theta}} + r \sin\theta d\phi \hat{\boldsymbol{\phi}}</math>; <math>d\tau = r^2 \sin\theta dr d\theta d\phi</math></p> <p><b>Gradient:</b> <math>\nabla f = \frac{\partial f}{\partial r} \hat{\mathbf{r}} + \frac{1}{r} \frac{\partial f}{\partial \theta} \hat{\boldsymbol{\theta}} + \frac{1}{r \sin\theta} \frac{\partial f}{\partial \phi} \hat{\boldsymbol{\phi}}</math></p> <p><b>Divergence:</b> <math>\nabla \cdot \mathbf{v} = \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 v_r) + \frac{1}{r \sin\theta} \frac{\partial}{\partial \theta} (\sin\theta v_\theta) + \frac{1}{r \sin\theta} \frac{\partial v_\phi}{\partial \phi}</math></p> <p><b>Curl:</b> <math>\nabla \times \mathbf{v} = \frac{1}{r \sin\theta} \left[ \frac{\partial}{\partial \theta} (\sin\theta v_\phi) - \frac{\partial v_\theta}{\partial \phi} \right] \hat{\mathbf{r}}</math>  <math>+ \frac{1}{r} \left[ \frac{1}{\sin\theta} \frac{\partial v_r}{\partial \phi} - \frac{\partial}{\partial r} (r v_\phi) \right] \hat{\boldsymbol{\theta}} + \frac{1}{r} \left[ \frac{\partial}{\partial r} (r v_\theta) - \frac{\partial v_r}{\partial \theta} \right] \hat{\boldsymbol{\phi}}</math></p> <p><b>Laplacian:</b> <math>\nabla^2 f = \frac{1}{r^2} \frac{\partial}{\partial r} \left( r^2 \frac{\partial f}{\partial r} \right) + \frac{1}{r^2 \sin\theta} \frac{\partial}{\partial \theta} \left( \sin\theta \frac{\partial f}{\partial \theta} \right) + \frac{1}{r^2 \sin^2\theta} \frac{\partial^2 f}{\partial \phi^2}</math></p> <p><b>Cylindrical.</b> <math>d\mathbf{l} = ds \hat{\mathbf{s}} + s d\phi \hat{\boldsymbol{\phi}} + dz \hat{\mathbf{z}}</math>; <math>d\tau = s ds d\phi dz</math></p> <p><b>Gradient:</b> <math>\nabla f = \frac{\partial f}{\partial s} \hat{\mathbf{s}} + \frac{1}{s} \frac{\partial f}{\partial \phi} \hat{\boldsymbol{\phi}} + \frac{\partial f}{\partial z} \hat{\mathbf{z}}</math></p> <p><b>Divergence:</b> <math>\nabla \cdot \mathbf{v} = \frac{1}{s} \frac{\partial}{\partial s} (s v_s) + \frac{1}{s} \frac{\partial v_\phi}{\partial \phi} + \frac{\partial v_z}{\partial z}</math></p> <p><b>Curl:</b> <math>\nabla \times \mathbf{v} = \left[ \frac{1}{s} \frac{\partial v_z}{\partial \phi} - \frac{\partial v_\phi}{\partial z} \right] \hat{\mathbf{s}} + \left[ \frac{\partial v_s}{\partial z} - \frac{\partial v_z}{\partial s} \right] \hat{\boldsymbol{\phi}} + \frac{1}{s} \left[ \frac{\partial}{\partial s} (s v_\phi) - \frac{\partial v_s}{\partial \phi} \right] \hat{\mathbf{z}}</math></p> <p><b>Laplacian:</b> <math>\nabla^2 f = \frac{1}{s} \frac{\partial}{\partial s} \left( s \frac{\partial f}{\partial s} \right) + \frac{1}{s^2} \frac{\partial^2 f}{\partial \phi^2} + \frac{\partial^2 f}{\partial z^2}</math></p>	<p><b>Triple Products</b></p> <p>(1) <math>\mathbf{A} \cdot (\mathbf{B} \times \mathbf{C}) = \mathbf{B} \cdot (\mathbf{C} \times \mathbf{A}) = \mathbf{C} \cdot (\mathbf{A} \times \mathbf{B})</math></p> <p>(2) <math>\mathbf{A} \times (\mathbf{B} \times \mathbf{C}) = \mathbf{B}(\mathbf{A} \cdot \mathbf{C}) - \mathbf{C}(\mathbf{A} \cdot \mathbf{B})</math></p> <p><b>Product Rules</b></p> <p>(3) <math>\nabla(fg) = f(\nabla g) + g(\nabla f)</math></p> <p>(4) <math>\nabla(\mathbf{A} \cdot \mathbf{B}) = \mathbf{A} \times (\nabla \times \mathbf{B}) + \mathbf{B} \times (\nabla \times \mathbf{A}) + (\mathbf{A} \cdot \nabla)\mathbf{B} + (\mathbf{B} \cdot \nabla)\mathbf{A}</math></p> <p>(5) <math>\nabla \cdot (f\mathbf{A}) = f(\nabla \cdot \mathbf{A}) + \mathbf{A} \cdot (\nabla f)</math></p> <p>(6) <math>\nabla \cdot (\mathbf{A} \times \mathbf{B}) = \mathbf{B} \cdot (\nabla \times \mathbf{A}) - \mathbf{A} \cdot (\nabla \times \mathbf{B})</math></p> <p>(7) <math>\nabla \times (f\mathbf{A}) = f(\nabla \times \mathbf{A}) - \mathbf{A} \times (\nabla f)</math></p> <p>(8) <math>\nabla \times (\mathbf{A} \times \mathbf{B}) = (\mathbf{B} \cdot \nabla)\mathbf{A} - (\mathbf{A} \cdot \nabla)\mathbf{B} + \mathbf{A}(\nabla \cdot \mathbf{B}) - \mathbf{B}(\nabla \cdot \mathbf{A})</math></p> <p><b>Second Derivatives</b></p> <p>(9) <math>\nabla \cdot (\nabla \times \mathbf{A}) = 0</math></p> <p>(10) <math>\nabla \times (\nabla f) = 0</math></p> <p>(11) <math>\nabla \times (\nabla \times \mathbf{A}) = \nabla(\nabla \cdot \mathbf{A}) - \nabla^2 \mathbf{A}</math></p> <p style="text-align: right;"><b>FUNDAMENTAL THEOREMS</b></p> <p><b>Gradient Theorem:</b> <math>\int_a^b (\nabla f) \cdot d\mathbf{l} = f(\mathbf{b}) - f(\mathbf{a})</math></p> <p><b>Divergence Theorem:</b> <math>\int (\nabla \cdot \mathbf{A}) d\tau = \oint \mathbf{A} \cdot d\mathbf{a}</math></p> <p><b>Curl Theorem:</b> <math>\int (\nabla \times \mathbf{A}) \cdot d\mathbf{a} = \oint \mathbf{A} \cdot d\mathbf{l}</math></p>
<p>Special case derivatives:                  (similar things true for <math>\mathcal{Z}</math>)</p>	
$\nabla \cdot \frac{\hat{\mathbf{r}}}{r^2} = 4\pi\delta(\mathbf{r})$	$\nabla^2 \frac{1}{r} = -4\pi\delta(\mathbf{r})$

BASIC EQUATIONS OF ELECTRODYNAMICS	FUNDAMENTAL CONSTANTS
<p><b>Maxwell's Equations</b></p> <p><i>In general:</i></p> $\begin{cases} \nabla \cdot \mathbf{E} = \frac{1}{\epsilon_0} \rho \\ \nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t} \\ \nabla \cdot \mathbf{B} = 0 \\ \nabla \times \mathbf{B} = \mu_0 \mathbf{J} + \mu_0 \epsilon_0 \frac{\partial \mathbf{E}}{\partial t} \end{cases}$ <p><i>In matter:</i></p> $\begin{cases} \nabla \cdot \mathbf{D} = \rho_f \\ \nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t} \\ \nabla \cdot \mathbf{B} = 0 \\ \nabla \times \mathbf{H} = \mathbf{J}_f + \frac{\partial \mathbf{D}}{\partial t} \end{cases}$	$\epsilon_0 = 8.85 \times 10^{-12} \text{ C}^2/\text{Nm}^2$ (permittivity of free space) $\mu_0 = 4\pi \times 10^{-7} \text{ N/A}^2$ (permeability of free space) $c = 3.00 \times 10^8 \text{ m/s}$ (speed of light) $e = 1.60 \times 10^{-19} \text{ C}$ (charge of the electron) $m = 9.11 \times 10^{-31} \text{ kg}$ (mass of the electron)
<p><b>Auxiliary Fields</b></p> <p><i>Definitions:</i></p> $\begin{cases} \mathbf{D} = \epsilon_0 \mathbf{E} + \mathbf{P} \\ \mathbf{H} = \frac{1}{\mu_0} \mathbf{B} - \mathbf{M} \end{cases}$ <p><i>Linear media:</i></p> $\begin{cases} \mathbf{P} = \epsilon_0 \chi_e \mathbf{E}, \quad \mathbf{D} = \epsilon \mathbf{E} \\ \mathbf{M} = \chi_m \mathbf{H}, \quad \mathbf{H} = \frac{1}{\mu} \mathbf{B} \end{cases}$	<p style="text-align: center;"><b>SPHERICAL AND CYLINDRICAL COORDINATES</b></p>
<p><b>Potentials</b></p> $\mathbf{E} = -\nabla V - \frac{\partial \mathbf{A}}{\partial t}, \quad \mathbf{B} = \nabla \times \mathbf{A}$	<p><b>Spherical</b></p> $\begin{cases} x = r \sin \theta \cos \phi \\ y = r \sin \theta \sin \phi \\ z = r \cos \theta \end{cases} \quad \begin{cases} \hat{\mathbf{x}} = \sin \theta \cos \phi \hat{\mathbf{r}} + \cos \theta \cos \phi \hat{\theta} - \sin \phi \hat{\phi} \\ \hat{\mathbf{y}} = \sin \theta \sin \phi \hat{\mathbf{r}} + \cos \theta \sin \phi \hat{\theta} + \cos \phi \hat{\phi} \\ \hat{\mathbf{z}} = \cos \theta \hat{\mathbf{r}} - \sin \theta \hat{\theta} \end{cases}$ $\begin{cases} r = \sqrt{x^2 + y^2 + z^2} \\ \theta = \tan^{-1}(\sqrt{x^2 + y^2}/z) \\ \phi = \tan^{-1}(y/x) \end{cases} \quad \begin{cases} \hat{\mathbf{r}} = \sin \theta \cos \phi \hat{\mathbf{x}} + \sin \theta \sin \phi \hat{\mathbf{y}} + \cos \theta \hat{\mathbf{z}} \\ \hat{\theta} = \cos \theta \cos \phi \hat{\mathbf{x}} + \cos \theta \sin \phi \hat{\mathbf{y}} - \sin \theta \hat{\mathbf{z}} \\ \hat{\phi} = -\sin \phi \hat{\mathbf{x}} + \cos \phi \hat{\mathbf{y}} \end{cases}$
<p><b>Lorentz force law</b></p> $\mathbf{F} = q(\mathbf{E} + \mathbf{v} \times \mathbf{B})$	<p><b>Cylindrical</b></p> $\begin{cases} x = s \cos \phi \\ y = s \sin \phi \\ z = z \end{cases} \quad \begin{cases} \hat{\mathbf{x}} = \cos \phi \hat{\mathbf{s}} - \sin \phi \hat{\phi} \\ \hat{\mathbf{y}} = \sin \phi \hat{\mathbf{s}} + \cos \phi \hat{\phi} \\ \hat{\mathbf{z}} = \hat{\mathbf{z}} \end{cases}$ $\begin{cases} s = \sqrt{x^2 + y^2} \\ \phi = \tan^{-1}(y/x) \\ z = z \end{cases} \quad \begin{cases} \hat{\mathbf{s}} = \cos \phi \hat{\mathbf{x}} + \sin \phi \hat{\mathbf{y}} \\ \hat{\phi} = -\sin \phi \hat{\mathbf{x}} + \cos \phi \hat{\mathbf{y}} \\ \hat{\mathbf{z}} = \hat{\mathbf{z}} \end{cases}$
<p><b>Energy, Momentum, and Power</b></p> <p><i>Energy:</i> <math>U = \frac{1}{2} \int (\epsilon_0 E^2 + \frac{1}{\mu_0} B^2) d\tau</math></p> <p><i>Momentum:</i> <math>\mathbf{P} = \epsilon_0 \int (\mathbf{E} \times \mathbf{B}) d\tau</math></p> <p><i>Poynting vector:</i> <math>\mathbf{S} = \frac{1}{\mu_0} (\mathbf{E} \times \mathbf{B})</math></p> <p><i>Larmor formula:</i> <math>P = \frac{\mu_0}{6\pi c} q^2 a^2</math></p>	

### Some miscellaneous mathematical stuff:

In[1]= (\* consider "rp" to mean "r prime" \*)

In[2]= Assuming[a > 0, Integrate[Exp[-2 rp/a], {rp, 0, r}]] // Expand

$$\text{Out[2]} = \frac{a}{2} - \frac{1}{2} a e^{-\frac{2r}{a}}$$

In[3]= Assuming[a > 0, Integrate[Exp[-2 rp/a], {rp, 0, Infinity}]] // Expand

$$\text{Out[3]} = \frac{a}{2}$$

In[4]= Assuming[a > 0, Integrate[Exp[-2 rp/a] rp, {rp, 0, r}]] // Expand

$$\text{Out[4]} = \frac{a^2}{4} - \frac{1}{4} a^2 e^{-\frac{2r}{a}} - \frac{1}{2} a e^{-\frac{2r}{a}} r$$

In[5]= Assuming[a > 0, Integrate[Exp[-2 rp/a] rp, {rp, 0, Infinity}]] // Expand

$$\text{Out[5]} = \frac{a^2}{4}$$

In[6]= Assuming[a > 0, Integrate[Exp[-2 rp/a] rp^2, {rp, 0, r}]] // Expand

$$\text{Out[6]} = \frac{a^3}{4} - \frac{1}{4} a^3 e^{-\frac{2r}{a}} - \frac{1}{2} a^2 e^{-\frac{2r}{a}} r - \frac{1}{2} a e^{-\frac{2r}{a}} r^2$$

In[7]= Assuming[a > 0, Integrate[Exp[-2 rp/a] rp^2, {rp, 0, Infinity}]] // Expand

$$\text{Out[7]} = \frac{a^3}{4}$$