

Bessel Functions, by Dr Colton Physics 442, Summer 2016

The Bessel functions, $J_\alpha(x)$ are a series of functions, that:


- (a) come up often, especially in partial differential equations
- (b) have interesting properties
- (c) are well understood and have been studied for centuries

They are typically only used for positive values of x . Here are the first four plotted.

Plots

BesselJ[α , x] is a built-in *Mathematica* function just like Sin[x], and gives $J_\alpha(x)$

```
In[25]= Plot[BesselJ[0, x], {x, 0, 20}]
```




Out[25]= - Graphics -

$J_0(x)$

crosses zero at 2.405, 5.520, 8.654, ...

← The only one that is not zero at the origin

```
In[26]= Plot[BesselJ[1, x], {x, 0, 20}]
```




Out[26]= - Graphics -

$J_1(x)$

crosses zero at 3.832, 7.016, 10.173, ...

```
In[27]= Plot[BesselJ[2, x], {x, 0, 20}]
```




Out[27]= - Graphics -

$J_2(x)$

crosses zero at 5.136, 8.417, 11.620, ...

```
In[28]= Plot[BesselJ[3, x], {x, 0, 20}]
```



Out[28]= - Graphics -

$J_3(x)$

crosses zero at 6.380, 9.761, 13.015, ...

Very important facts

- Can be computed via a series formula: $J_\alpha(x) = \sum_{k=0}^{\infty} \frac{(-1)^k x^{2k+\alpha}}{k!(k+\alpha)!2^{2k+\alpha}}$.
- Derivatives:
 - $\frac{d}{dx}(J_0(x)) = -J_1(x)$
 - $\frac{d}{dx}(J_n(x)) = \frac{J_{n-1}(x) - J_{n+1}(x)}{2}$
- Zeroes: $u_{\alpha m}$ represents the m^{th} zero of $J_\alpha(x)$. From the previous page:
 - $u_{01} = 2.405, u_{02} = 5.520, u_{03} = 8.654, \dots$
 - $u_{11} = 3.832, u_{12} = 7.016, u_{13} = 10.173, \dots$
 - etc.


These numbers are available in Mathematica via the BesselJZero function: BesselJZero[0,1] yields a result of 2.4048255576957727686..., etc.

- Differential equations (“Bessel’s equation”):
 - $x^2 \frac{d^2 f}{dx^2} + x \frac{df}{dx} + (x^2 - \alpha^2)f = 0$ has solution $J_\alpha(x)$, or linear combinations
 - $x^2 \frac{d^2 f}{dx^2} + x \frac{df}{dx} + (u_{\alpha m}^2 x^2 - \alpha^2)f = 0$ has solution $J_\alpha(u_{\alpha m} x)$, or linear combinations
- There are a second set of solutions to Bessel’s differential equation, called $Y_\alpha(x)$ (sometimes written $N_\alpha(x)$) or the “Bessel functions of the second kind”; these diverge at $x = 0$ and are therefore often not used.
 - Various other linear combinations of $J_\alpha(x)$ and $Y_\alpha(x)$ are sometimes used; two examples are the “modified Bessel functions” and the “Hankel functions”. These are beyond the scope of this course.
 - The so-called “spherical Bessel functions” (and “spherical Hankel functions”) are solutions to a different, albeit closely related, differential equation. These are also beyond the scope of this course.
- When $u_{\alpha m}$ is added to the argument in the form $u_{\alpha m} r$, the functions go to zero at $r = 1$. Then e.g. $J_0(u_{0m} r)$ represents a series of functions between the origin and a circular boundary of $r = 1$, having m antinodes. For example $J_0(u_{02} x)$ has two antinodes, $J_0(u_{03} x)$ has three antinodes, etc.

```

In[9] = u02 = BesselJZero[0, 2];
Plot3D[BesselJ[0, u02 Sqrt[x^2 + y^2]], {x, -1, 1}, {y, -1, 1},
RegionFunction -> Function[{x, y, z}, x^2 + y^2 < 1]]


```



```

In[10] = u03 = BesselJZero[0, 3];
Plot3D[BesselJ[0, u03 Sqrt[x^2 + y^2]], {x, -1, 1}, {y, -1, 1},
RegionFunction -> Function[{x, y, z}, x^2 + y^2 < 1]]

```



- Orthogonality: $J_0(u_{0m} x)$ and $J_0(u_{0n} x)$ are orthogonal over the interval $(0, 1)$ with respect to a weighting function of x :

$$\int_0^1 x J_0(u_{0m} x) J_0(u_{0n} x) dx = \begin{cases} 0 & \text{if } n \neq m \\ \frac{1}{2} (J_1(u_{0m} x))^2 & \text{if } n = m \end{cases}$$


Or, more generally for $J_\alpha(x)$:

$$\int_0^1 x J_\alpha(u_{\alpha m} x) J_\alpha(u_{\alpha n} x) dx = \begin{cases} 0 & \text{if } n \neq m \\ \frac{1}{2} (J_{\alpha+1}(u_{\alpha m} x))^2 & \text{if } n = m \end{cases}$$


Orthogonality, depicted

```
In[1]:= u02 = BesselJZero[0, 2];
        u03 = BesselJZero[0, 3];
```


```
In[3]:= Plot[BesselJ[0, u02 x], {x, 0, 3}]
```



```
In[4]:= Plot[BesselJ[0, u03 x], {x, 0, 3}, PlotRange -> All]
```




```
In[5]:= Plot[x BesselJ[0, u02 x] BesselJ[0, u03 x], {x, 0, 1}, Filling -> Axis]
```



```
In[6]:= Integrate[x BesselJ[0, u02 x] BesselJ[0, u03 x], {x, 0, 1}]
```

```
Out[6]:= 0
```

```
In[7]:= Plot[x BesselJ[0, u03 x] BesselJ[0, u03 x], {x, 0, 1}, Filling -> Axis]
```



```
In[8]:= Integrate[x BesselJ[0, u03 x] BesselJ[0, u03 x], {x, 0, 1}] // N
```

```
Out[8]:= 0.0368432
```

```
In[9]:= .5 (BesselJ[1, u03])^2 // N
```

```
Out[9]:= 0.0368432
```

These are two functions which are orthogonal. They are both J_0 functions, but one has been scaled so that its *second* “zero crossing” occurs at $x = 1$, and the other has been scaled so that its *third* zero crossing occurs at $x = 1$.

The two functions are only orthogonal over the range from 0 to 1, and only with a weighting function of x . But within those constraints, you can see that the positive areas cancel out the negative areas.

The integral is zero.

$$\int_0^1 x J_0(u_{0m}x) J_0(u_{0n}x) dx$$

equals 0 if $n \neq m$!

By contrast, $J_0(u_{0m}x)$ is not orthogonal to itself.

$$\int_0^1 x J_0(u_{0m}x) J_0(u_{0n}x) dx$$

equals $\frac{1}{2} (J_1(u_{0m}x))^2$ if $n = m$!

Comparison with sines & cosines:

Sines/Cosines

- Two oscillatory functions: $\sin(x)$ and $\cos(x)$. Sometimes one of them is not used, due to the symmetry of the problem.
- You determine the value of $\sin(x)$ or $\cos(x)$ for arbitrary x by using a calculator or computer program.

$$\sin(x) = \sum_{k=0}^{\infty} \frac{(-1)^k x^{2k+1}}{(2k+1)!}$$

$$\cos(x) = \sum_{k=0}^{\infty} \frac{(-1)^k x^{2k}}{(2k)!}$$

Consider just $\sin(x)$:

- The zeroes of $\sin(x)$ are at $x = \pi, 2\pi, 3\pi$, etc.
 $x = "m\pi"$ is the m^{th} zero
- Using the zeroes in the argument, $\sin(m\pi x)$ has m antinodes in the interval from 0 to 1. And at $x = 1$ (the boundary), $\sin(m\pi x) = 0$ for all m .
- The differential equation satisfied by $f = \sin(x)$ is $f'' + f = 0$.

The differential equation satisfied by $f = \sin(m\pi x)$ is $f'' + (m\pi)^2 f = 0$.

- $\sin(n\pi x)$ is orthogonal to $\sin(m\pi x)$ on the interval $(0,1)$:

$$\int_0^1 \sin(n\pi x) \sin(m\pi x) dx = \begin{cases} 0, & \text{if } n \neq m \\ \frac{1}{2}, & \text{if } n = m \end{cases}$$

Bessel functions

Two oscillatory functions for each α : $J_\alpha(x)$ and $Y_\alpha(x)$. Typically Y_α is not used because it's infinite at the origin.

You determine the value of $J_\alpha(x)$ for arbitrary x by using a calculator or computer program.

$$J_\alpha(x) = \sum_{k=0}^{\infty} \frac{(-1)^k x^{2k+\alpha}}{k!(k+\alpha)! 2^{2k+\alpha}}$$

Consider just $J_\alpha(x)$ for one α , say $\alpha = 0$:

(similar things hold true for all α 's)

The zeroes of $J_0(x)$ are at $x \approx 2.405, 5.520, 8.654$, etc.

$x = "u_{0m}"$ is the m^{th} zero

Using the zeroes in the argument, $J_0(u_{0m}r)$ has m antinodes in the circular region $r \leq 1$. And at $r = 1$ (the boundary), $J_0(u_{0m}x) = 0$ for all m .

The differential equation satisfied by $f = J_0(x)$ is $x^2 f'' + x f' + (x^2 - 0^2) f = 0$.

The differential equation satisfied by $f = J_0(u_{0m}x)$ is $x^2 f'' + x f' + (u_{0m}^2 x^2 - 0^2) f = 0$.

$0^2 \rightarrow \alpha^2$ for other α 's

$J_0(u_{0n}x)$ is orthogonal to $J_0(u_{0m}x)$ on the interval $(0,1)$, with respect to a weighting function of x :

$$\int_0^1 x J_0(u_{0n}x) J_0(u_{0m}x) dx = \begin{cases} 0, & \text{if } n \neq m \\ \frac{1}{2} (J_1(u_{0m}))^2, & \text{if } n = m \end{cases}$$

Similar orthogonality holds for other values of α .

Additionally, the Bessel functions are related to sines/cosines through this integral formula:

$$J_\alpha(x) = \frac{1}{\pi} \int_0^\pi \cos(\alpha\theta - x \sin \theta) d\theta$$

Quote from Mary Boas, in *Mathematica Methods in the Physical Sciences*: "In fact, if you had first learned about $\sin(nx)$ and $\cos(nx)$ as power series solutions of $y'' = -n^2 y$, instead of in elementary trigonometry, you would not feel that Bessel functions were appreciably more difficult or strange than trigonometric functions. Like sines and cosines, Bessel functions are solutions of a differential equation; they are tabulated and their graphs can be drawn; they can be represented as a series; and a large number of formulas about them are known."